

Confid

(a) contacting said nucleic acid sequence, in one amplification reaction or separate amplification reactions, with a first pair of PCR primers and a second pair of PCR primers under conditions that allow hybridization of said PCR primers to said nucleic acid sequence, a first member of said first pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said first pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said first pair hybridizes, and a first member of said second pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said second pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said second pair hybridizes, said PCR primers being characterized as follows:

Q1

(i) one of said first pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to a first allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to a second allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at a single nucleotide that is disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said first pair of primers is capable of amplifying said first allele under appropriate conditions; and

(ii) one of said second pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at one or more

a1  
a1  
nucleotides that are disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said second pair of primers is capable of amplifying said first allele under appropriate conditions;

(b) carrying out said amplification reaction or reactions; and

(c) detecting any amplification product of step (b), wherein the presence of amplification product is indicative of the presence of said first allele in said nucleic acid sequence.

---

6. (Amended) The method of claim 1, wherein said one of said second pair of PCR primers in step (a)(ii) includes at least two non-complementary nucleotides that are disposed within the five nucleotides adjacent to the 3'-terminal nucleotide of each primer.

a2  
7. (Amended) The method of claim 1, wherein said polymorphic sequence comprises a single nucleotide polymorphism.

8. (Amended) The method of claim 1, wherein said one of said first pair of PCR primers in step (a)(i) and said one of said second pair of PCR primers in step (a)(ii) also comprise a universal primer binding site.

a3  
10. (Amended) The method of claim 1, wherein said one of said first pair of PCR primers in step (a)(i) and said one of said second pair of PCR primers in step (a)(ii) also comprise a unique hybridization tag.

a4 12. (Amended) The method of claim 10, wherein said detection step is carried

out on a solid support to which a binding partner for each hybridization tag is immobilized.

14. (Amended) The method of claim 1, further comprising:

as (d) contacting said nucleic acid sequence, in one amplification reaction or separate amplification reactions, with a third pair of PCR primers and a fourth pair of PCR primers under conditions that allow hybridization of said PCR primers to said nucleic acid sequence, a first member of said third pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said third pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said third pair hybridizes, and a first member of said fourth pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said fourth pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said fourth pair hybridizes, said PCR primers being characterized as follows:

(i) one of said third pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at a single nucleotide

that is disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said third pair of primers is capable of amplifying said second allele under appropriate conditions; and

(ii) one of said fourth pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at one or more nucleotides that are disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said fourth pair of primers is capable of amplifying said second allele under appropriate conditions; and

(e) carrying out said amplification reaction or reactions; and

(f) detecting any amplification product of step (e), wherein the presence of amplification product is indicative of the presence of said second allele in said nucleic acid sequence.

15. (Amended) A kit for determining whether a nucleic acid sequence comprises a particular allele of a polymorphic sequence, said kit comprising:

(a) a first pair of PCR primers and a second pair of PCR primers, a first member of said first pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said first pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said first pair hybridizes, and a first member

of said second pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said second pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said second pair hybridizes, wherein said PCR primers are characterized as follows:

cont'd  
(i) one of said first pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to a first allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to a second allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at a single nucleotide that is disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said first pair of primers is capable of amplifying said first allele under appropriate conditions; and

AS  
(ii) one of said second pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at one or more nucleotides that are disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said second pair of primers is capable of amplifying said first allele under appropriate conditions.

16. (Amended) The kit of claim 15, further comprising:

(b) a third pair of PCR primers and a fourth pair of PCR primers, a first member of said third pair of PCR primers hybridizing to one strand of said nucleic acid

Cont'd  
OS  
sequence and the other member of said third pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said third pair hybridizes, and a first member of said fourth pair of PCR primers hybridizing to one strand of said nucleic acid sequence and the other member of said fourth pair of PCR primers hybridizing to the other strand of said nucleic acid sequence on the opposite side of said polymorphic sequence from the side to which said first member of said fourth pair hybridizes, wherein said PCR primers are characterized as follows:

(i) one of said third pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at a single nucleotide that is disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said third pair of primers is capable of amplifying said second allele under appropriate conditions; and

(ii) one of said fourth pair of PCR primers is (a) complementary at its 3'-terminal nucleotide to said second allele of said polymorphic sequence, (b) non-complementary at its 3'-terminal nucleotide to said first allele of said polymorphic sequence, and (c) non-complementary to said nucleic acid sequence at one or more nucleotides that is disposed within the five nucleotides adjacent to said 3'-terminal nucleotide, wherein said fourth pair of primers is capable of amplifying said second allele under appropriate conditions.